

Guacamayos for some exceptional studies.

ORANGE-BELLIED EUPHONIA (*Euphonia xanthogaster*) – Probably the most common euphonia in Ecuador as it is so wide ranging and able to tolerate many different habitat types.

BLUE-NAPED CHLOROPHONIA (*Chlorophonia cyanea longipennis*) – A heard bird for just about everybody, but Lois managed to spot one with a flock - where we were hearing them - on the S. slope of the Guacamayos.

YELLOW-BELLIED SISKIN (*Spinus xanthogastrus*) – Nice looks at this western foothill species around Milpe.

HOODED SISKIN (*Spinus magellanicus capitalis*) – Common in the central valley; we saw this brightly colored species on our first day.

OLIVACEOUS SISKIN (*Spinus olivaceus*) – Brief views in the east.

MAMMALS

WHITE-EARED OPOSSUM (*Didelphis albiventris*) – Quick views on our last full day at San Isidro.

BRAZILIAN RABBIT (*Sylvilagus brasiliensis*) – A fairly common mammal that is mostly encountered in the highland grasslands, and we saw them a couple of times in such habitats.

RED-TAILED SQUIRREL (*Sciurus granatensis*) – Common on both slopes.

BLACK AGOUTI (*Dasyprocta fuliginosa*) – Seen around the gardens at San Isidro, where they frequently forage about near the cabins and dining room.

ADDITIONAL COMMENTS

Totals for the tour: 342 bird taxa and 4 mammal taxa